

Weakly coupled two-slow–two-fast systems, folded singularities and mixed mode oscillations

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2014 Nonlinearity 27 1555

(<http://iopscience.iop.org/0951-7715/27/7/1555>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 129.187.254.47

This content was downloaded on 29/06/2014 at 10:12

Please note that [terms and conditions apply](#).

Weakly coupled two-slow–two-fast systems, folded singularities and mixed mode oscillations

M Krupa¹, B Ambrosio and M A Aziz-Alaoui²

¹ INRIA Paris-Rocquencourt Research Center, Domaine de Voluceau BP 105, 78153 Le Chesnay cedex, France

² Normandie Univ, France; ULH, LMAH, F-76600 Le Havre; FR CNRS 3335, 25 rue Philippe Lebon 76600 Le Havre, France

E-mail: benjamin.ambrosio@univ-lehavre.fr

Received 27 January 2013, revised 28 October 2013

Accepted for publication 30 January 2014

Published 24 June 2014

Recommended by D V Treschev

Abstract

We study mixed mode oscillations (MMOs) in systems of two weakly coupled slow/fast oscillators. We focus on the existence and properties of a folded singularity, called folded saddle node of type II (FSN II), that allows the emergence of MMOs in the presence of a suitable global return mechanism. As FSN II corresponds to a transcritical bifurcation for a desingularized reduced system, we prove that, under certain non-degeneracy conditions, such a transcritical bifurcation exists. We then apply this result to the case of two coupled systems of FitzHugh–Nagumo type. This leads to a non-trivial condition on the coupling that enables the existence of MMOs.

Keywords: slow–fast dynamics, mixed mode oscillations, transcritical bifurcation, folded saddle-node, canard

Mathematics Subject Classification: 34C15, 34D15, 34K18

(Some figures may appear in colour only in the online journal)

1. Introduction

The *Canard phenomenon*, known also as the *canard explosion* is a transition from a small, Hopf type oscillation to a relaxation oscillation, occurring upon variation of a parameter. This transition was first found in the context of the van der Pol equation [2], and soon after in numerous models of phenomena occurring in engineering and in chemical reactions [4].

A common feature of all these models is the presence of time scale separation (one slow, one fast variable). A particular feature of canard explosion is that it takes place in a very small parameter interval. For the van der Pol system, where the ratio of the timescales is given by a small parameter ε , the width of this parameter interval can be estimated by $O(\exp(-c/\varepsilon))$, where $c > 0$ is a fixed positive constant³. The transition occurs through a sequence of *canard cycles*, whose striking feature is that they contain segments following closely a stable slow manifold and subsequently an unstable slow manifold.

The work on canard explosion led to investigations of slow/fast systems in three dimensions, with two slow and one fast variables. In the context of these systems, the concept of a *canard solution* or simply *canard* has been introduced as a solution passing from a stable to an unstable slow manifold [1, 10, 13]. Canards arise near so called folded singularities, of which the best known is folded node [1, 13] and [15]. Unlike in systems with one slow variable, canards occur robustly (in $O(1)$ parameter regions) in systems with two slow variables. Related to canards are mixed mode oscillations (MMOs), which are trajectories that combine small oscillations and large oscillations of relaxation type, both recurring in an alternating manner. Recently, there has been a lot of interest in MMOs that arise due to the presence of canards, starting with the work of Milik *et al* [10]. The small oscillations arise during the passage of the trajectories near a folded node or a related folded singularity. The dynamics near the folded singularity is transient, yet recurrent: the trajectories return to the neighbourhood of the folded singularity by way of a global return mechanism [3].

The setting of a folded node combined with a global return mechanism, elucidated in [3], has led to the explanations of MMO dynamics found in applications [7, 11, 12, 14]. A shortcoming of the folded node setting is the lack of connection to a Hopf bifurcation, which seems to play a prominent role in many MMOs. This led to the interest in another, more degenerate, folded singularity, known as *folded saddle node of type II* (FSN II), originally introduced in [10] and recently analyzed in some detail by Krupa and Wechselberger [9]. Guckenheimer [8] studied a very similar problem in the parameter regime, which was yet closer to the Hopf bifurcation, calling it *singular Hopf bifurcation*. The transition between the two settings was studied in [5]. Another interesting singularity, which was mentioned in [9] and can lead to rich families of MMOs, is the *folded saddle node of type I* (FSN I). In the case of this bifurcation, small oscillations seem to be related to the presence of a delayed Hopf bifurcation rather than a true Hopf bifurcation. For a more comprehensive overview we refer the reader to the recent review paper [6]. The theory of two slow and one fast variables has recently been generalized by Wechselberger [17] to arbitrary finite dimensions.

In this paper, we study systems of two weakly coupled slow/fast oscillators. We assume that in the absence of the coupling one of the oscillators is undergoing a canard explosion and the other is in general position. We show that turning on the weak coupling leads to the occurrence of MMOs. We focus on the very interesting case of FSN II, which, in the uncoupled case, corresponds to one of the oscillators undergoing a canard explosion while the other is at a stable equilibrium. As elaborated in [3], canard induced MMOs arise through a combined presence of a folded singularity and suitable return mechanism. In this paper we focus on the existence and properties of a folded singularity, leaving the study of the return mechanism for future investigations.

This paper is organized as follows. We start with a background section, section 2, in which we explain the very standard case of MMOs in the context of two slow and one fast variables and, subsequently, the case of two slow and two fast variables with a simple fold curve. This

³ Strictly speaking, if one defines canard explosion transition between small canards and canard cycles with a large head, then the width of the parameter interval where the transition takes place is given by $O(\exp(-c/\varepsilon))$.

material is included for completeness and presented in such a way that the coupled oscillator case becomes a simple corollary. In section 3, we treat the general case of coupled oscillators and in section 4, we consider the example of two coupled FitzHugh–Nagumo systems.

2. Background

2.1. The basic case of one fast and two slow variables

We consider a system of the form

$$\epsilon \dot{x} = f(x, y, \epsilon), \tag{1}$$

$$\dot{y} = g(x, y, \epsilon), \quad x \in \mathbb{R}, \quad y = (y_1, y_2) \in \mathbb{R}^2, \quad g = (g_1, g_2). \tag{2}$$

The associated reduced system is

$$0 = f(x, y, 0), \tag{3}$$

$$\dot{y} = g(x, y, 0), \quad x \in \mathbb{R}, \quad y = (y_1, y_2) \in \mathbb{R}^2. \tag{4}$$

Let S_0 denote the surface defined by $0 = f(x, y, 0)$. Non-hyperbolic points correspond to the points on S_0 , for which $f_x = 0$ (we use the notation $f_\xi = \partial f / \partial \xi$). At such points, the equation $0 = f(x, y, 0)$ cannot be solved for x as a function of y . Suppose $(0, 0)$ is a non-hyperbolic point. We make a non-degeneracy assumption $f_{xx}(0) \neq 0$. In order to obtain an explicit equation for the slow flow, we try to solve $0 = f(x, y, 0)$ for y_1 or y_2 . It is convenient to first change the variables to simplify the process of finding such a solution. We begin with a change of variables of the form $x \rightarrow x + \eta(y)$, $\eta(0, 0) = 0$, where v satisfies $f_x(v(y), y) = 0$. In the new variables, with additional scaling, $f(x, y, 0)$ has the form

$$f(x, y_1, y_2) = f(0, y, 0) + x^2 + O(x^2|y|, x^3).$$

In this last equation, and in the following, we do not write the variable ϵ in the function f , which is always equal to 0; that is, we write $f(x, y_1, y_2)$ for $f(x, y_1, y_2, 0)$. We make a non-degeneracy assumption $f_y(0, 0) \neq (0, 0)$. Without loss of generality (WLOG), we can assume that $f_{y_1}(0, 0) \neq 0$. We can now introduce a new coordinate $\tilde{y}_1 = -f(0, y)$ and immediately drop the tilde to simplify the notation. In the new coordinates, f has the form

$$f(x, y_1, y_2) = -y_1 + x^2 + O(x^2|y|, x^3).$$

The transformations we made may be just valid locally; that is, only in a small neighbourhood of the non-hyperbolic point. Now S_0 is (locally) represented as a graph:

$$y_1 = x^2(1 + O(y_2, x)). \tag{5}$$

and the fold is the straight line $x = y_1 = 0$.

We now transform (3)–(4), first removing the constraint $f(x, y) = 0$ and subsequently desingularising the resulting equation. Differentiating (5) we get

$$\dot{y}_1 = 2x(1 + O(y_2, x))\dot{x} + O(x^2). \tag{6}$$

Substituting (6) into (4) we get

$$2x(1 + O(y_2, x))\dot{x} = g_1(x, x^2(1 + O(y_2, x)), y_2) + O(x^2) \tag{7}$$

$$\dot{y}_2 = g_2(x, x^2(1 + O(y_2, x)), y_2). \tag{8}$$

Now it is clear that (7) is singular at the fold as long as $g_1(0, 0, y_2) \neq 0$. Points on the fold line for which $g_1(0, 0, y_2) = 0$ are called folded singularities. We would like to understand the flow near such points better and to this end we apply a singular time rescaling to (7)–(8)

by multiplying the right-hand side (rhs) by the factor $2x(1 + O(y_2, x))$ and cancelling it in (7). This leads to the following system:

$$\dot{x} = g_1(x, x^2(1 + O(y_2, x)), y_2) + O(x^2) \quad (9)$$

$$\dot{y}_2 = 2x(1 + O(y_2, x))g_2(x, x^2(1 + O(y_2, x)), y_2). \quad (10)$$

We refer to (9)–(10) as the desingularized system. Note that folded singularities correspond to equilibria of (9)–(10) with $x = 0$. Further note that the trajectories of (7)–(8) and (9)–(10) restricted to the half plane $x > 0$ differ only by time parametrization but are the same as sets. The trajectories in the half-plane $x < 0$, on the attracting part of the critical manifold, are the same as sets but have the opposite direction of time. The flow near folded singularities is determined by the linearization of (9)–(10) at folded singularities, which is given by the Jacobian matrix:

$$\begin{pmatrix} g_{1,x}(0, 0, y_2) & g_{1,y_2}(0, 0, y_2) \\ 2g_2(0, 0, y_2) & 0 \end{pmatrix}. \quad (11)$$

Folded singularities are classified according to the type of the corresponding equilibrium of (9)–(10). Canards arise near folded saddles, and folded nodes and small oscillations are associated with folded nodes, see the work of Benoît [1], and Szmolyan and Wechselberger [13]. A folded node, which is the only generic folded singularity whose dynamics is accompanied by small oscillations, occurs when the corresponding equilibrium of (9)–(10) has two real positive eigenvalues (recall that we have changed the direction of the flow on the attracting side of the critical manifold).

2.2. Folded singularities

We now consider the case when (f, g) (the rhs of (1)–(2)) depends on a regular parameter and $g_2(0, 0, y_2)$ passes through 0 when this parameter is varied. Suppose that the equation $g_2(0, 0, y_2) = 0$ admits a unique solution \bar{y}_2 . If we assume that $g_{1,y_2}(0, 0, y_2) \neq 0$ for y_2 in a neighborhood of \bar{y}_2 and $\frac{\partial}{\partial y_2} g_2(0, 0, \bar{y}_2) \neq 0$, then this passage corresponds to a change of sign of the determinant of the Jacobian (11), from negative to positive, or vice versa. We make the assumption that $g_{1,x} > 0$, which guarantees that the flow of (7)–(8), on the stable part of the critical manifold is towards the fold. This, in turn, means that the eigenvalues change from one positive and one negative (negative determinant) to two positive (positive determinant); that is, from saddle to node. It follows that this transition corresponds to the onset of small oscillations. Hence, the onset of small oscillations is a consequence of the passage of $g_2(0, 0, y_2)$ through 0. This transition is called FSN II.

Since finding FSN II in coupled oscillator systems is the focus of this paper, we review some of the features of this bifurcation, referring the reader to [9] for details. An important feature of FSN II is that it corresponds to the passage of a true equilibrium of (3)–(4) through the fold line. We assume WLOG that FSN II corresponds to the point $(x, y, \lambda) = (0, 0, 0)$, where λ is the regular parameter. Suppose that FSN II is *non-degenerate*, that is $(x, y, \lambda) = (0, 0, 0)$ corresponds to a transcritical bifurcation of (9)–(10). Then, for $\lambda = 0$, there is a folded singularity satisfying $g_2(0, 0, 0) = 0$ and, for $\lambda \neq 0$ but close to 0, there is a point close to the origin, such that $g_1(x, x^2 + O(x^3), y_2) = g_2(x, x^2 + O(x^3), y_2) = 0$; that is, a true equilibrium of (3)–(4) near the fold line. For the original system (1)–(2) one can prove that this corresponds to a so called singular Hopf bifurcation [8], and, in a different regime of the parameter λ , to a delayed Hopf bifurcation [9]. These two bifurcations lead to the existence of small oscillations and thus enable the existence of MMOs. We note here that the interesting case of FSN II is when the equilibrium is stable on the stable slow manifold and a saddle on the unstable slow manifold.

For folded nodes, which are not close to either FSN II or to FSN I, defined by $g_{1,y_2} = 0$, small oscillations are extremely small, which means that they cannot be seen, except with very detailed numerics, see [3]. Hence, the most interesting cases occur near FSN I or FSN II.

2.3. Canonical system in two fast and two slow dimensions

We consider a system with two fast and two slow variables

$$\varepsilon \dot{x} = f(x, y, \varepsilon), \tag{12}$$

$$\dot{y} = g(x, y, \varepsilon), \quad x = (x_1, x_2) \in \mathbb{R}^2, \quad y = (y_1, y_2) \in \mathbb{R}^2, \tag{13}$$

$f = (f_1, f_2), g = (g_1, g_2)$. Recall that in the fast formulation (12)–(13) has the form

$$x' = f(x, y, \varepsilon), \tag{14}$$

$$y' = \varepsilon g(x, y, \varepsilon). \tag{15}$$

Recall also that the reduced problem has the form

$$0 = f(x, y, 0), \tag{16}$$

$$\dot{y} = g(x, y, 0). \tag{17}$$

and the layer problem has the form

$$x' = f(x, y, 0), \tag{18}$$

$$y' = 0. \tag{19}$$

Let $\Phi : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ be the map defined by

$$\Phi(x, y) = \begin{pmatrix} f_1(x, y) \\ f_2(x, y) \\ \det(f_x)(x, y) \end{pmatrix}. \tag{20}$$

The fold curve is defined by the condition $\Phi = 0$. We consider a point on the fold curve (WLOG we assume that this point is the origin $(0, 0)$). We now state a few conditions which assure that the fold curve is *simple*; naturally the first condition is that the linearization of layer system has a simple eigenvalue 0. More specifically, our first assumption is as follows:

$$f_x(0, 0) \text{ has one simple } 0 \text{ eigenvalue and one simple eigenvalue in the left half-plane.} \tag{21}$$

Recall that in section 2.1 we made additional assumptions, namely $f_{xx}(0, 0) \neq 0$ and $f_y(0, 0) \neq (0, 0)$. Here, we generalize these conditions in the following way. Let (v_1, v_2) denote the null vector (the eigenvector of 0). Our additional non-degeneracy conditions is as follows:

$$D\Phi(0, 0) \text{ has full rank and } \kappa = (\det(f_x))_x(0, 0) \cdot (v_1, v_2) \neq 0. \tag{22}$$

We have the following lemma.

Lemma 1. Hypothesis (22) implies that either $\Phi_{(x_1, x_2, y_1)}(0, 0)$ or $\Phi_{(x_1, x_2, y_2)}(0, 0)$ is invertible.

Proof. Let (w_1, w_2) be the eigenvector of the negative eigenvalue. Note that either $f_{y_1}(0, 0)$ or $f_{y_2}(0, 0)$ must be linearly independent of (w_1, w_2) , otherwise $D\Phi(0, 0)\mathbb{R}^4$ would be in the span of $(w_1, w_2, 0)$ and $(0, 0, 1)$. WLOG we assume that $f_{y_1}(0, 0)$ is independent of (w_1, w_2) and let $M = \Phi_{(x_1, x_2, y_1)}(0, 0)$. Note that $M(v_1, v_2, 0) = (0, 0, \kappa)$ and that the first two coordinates of $M(w_1, w_2, 0)$ are a multiple of (w_1, w_2) . Note also that the first two coordinates of $M(0, 0, 1)$ are equal to $f_{y_1}(0, 0)$. Since $f_{y_1}(0, 0)$ is independent of (w_1, w_2) , the vectors $M(v_1, v_2, 0), M(w_1, w_2, 0)$ and $M(0, 0, 1)$ are linearly independent.

We now have the following proposition.

Proposition 1. *We assume that $(0, 0)$ is on the fold curve (i.e. $\Phi(0, 0) = 0$) and that conditions (21)–(22) are satisfied. Then, there exists a neighbourhood $U = V \times W$ of $(0, 0) \in \mathbb{R}^4$, and a unique function $\psi : V \subset \mathbb{R} \rightarrow \mathbb{R}^3$, such that:*

$$\{(x, y) \in U; \Phi = 0\} = \{(x, y_1) \in \mathbb{R}^3; (x, y_1) = \psi(y_2), y_2 \in V\}.$$

Proof. By lemma (1) the matrix $M = \Phi_{(x_1, x_2, y_1)}(0, 0)$ is non singular. Hence, the equation $\Phi = 0$ can be solved by the implicit function theorem, giving (x_1, x_2, y_1) as a function of y_2 .

In the remainder of this section we will assume that the fold curve has been transformed to the coordinate line $(0, 0, 0, y_2)$ and that the fast system has been diagonalized. More specifically, we assume that f satisfies the following conditions

$$f(0, 0, 0, y_2, 0) = 0, \quad \frac{\partial f_1}{\partial x_1}(0, 0, 0, y_2, 0) = 0, \quad \frac{\partial f_2}{\partial x_2}(0, 0, 0, y_2, 0) = \lambda(y_2) < 0, \quad (23)$$

$$\frac{\partial f_1}{\partial x_2}(0, 0, 0, y_2, 0) = \frac{\partial f_2}{\partial x_1}(0, 0, 0, y_2, 0) = 0. \quad (24)$$

Note that the non-degeneracy condition (22) reduces to

$$\frac{\partial f_1}{\partial y_1}(0, 0, 0, y_2, 0) \neq 0, \quad \frac{\partial^2 f_1}{\partial x_1^2}(0, 0, 0, y_2, 0) \neq 0. \quad (25)$$

WLOG we assume

$$\frac{\partial f_1}{\partial y_1}(0, 0, 0, y_2, 0) = -1, \quad \frac{\partial^2 f_1}{\partial x_1^2}(0, 0, 0, y_2, 0) = 2.$$

We can now expand f in Taylor series:

$$f(x, y, \varepsilon) = \begin{pmatrix} -y_1 + x_1^2 + O(x_1 x_2, x_2^2, x_1 y_1, x_2 y_1, y_1^2) + O(\|(x, y)\|^3) + O(\varepsilon) \\ \lambda(y)x_2 + O(y_1, \|x\|^2) + O(\varepsilon) \end{pmatrix}. \quad (26)$$

The defining conditions of the slow manifold are $f(x, y, 0) = 0$, or

$$0 = -y_1 + x_1^2 + O(x_1 x_2, x_2^2, x_1 y_1, x_2 y_1, y_1^2) + O(\|(x, y)\|^3), \quad (27)$$

$$0 = \lambda(y)x_2 + O(y_1, \|x\|^2). \quad (28)$$

From (28) we get, by the implicit function theorem, $x_2 = p(x_1, y)$, with $p(x_1, y) = O(y_1, x_1^2)$. Plugging into (27) we get the usual condition $y_1 = x_1^2 + O(x_1^3)$. Following the approach of section 2.1 we substitute $y_1 = x_1^2 + O(x_1^3)$ into (17) getting

$$(2x_1 + O(x_1^2))\dot{x}_1 = g_1(x_1, p(x_1, x_1^2 + O(x_1^3)), x_1^2 + O(x_1^3), y_2, 0), \quad (29)$$

$$\dot{y}_2 = g_2(x_1, p(x_1, x_1^2 + O(x_1^3)), y_2), x_1^2 + O(x_1^3), y_2, 0). \quad (30)$$

Finally, we get the desingularized equation as follows:

$$\dot{x}_1 = g_1(x_1, p(x_1, x_1^2 + O(x_1^3)), y_2), x_1^2 + O(x_1^3), y_2, 0), \quad (31)$$

$$\dot{y}_2 = g_2(x_1, p(x_1, x_1^2 + O(x_1^3)), y_2), x_1^2 + O(x_1^3), y_2, 0)(2x_1 + O(x_1^2)). \quad (32)$$

Folded singularities are points $(0, y_2)$ with y_2 satisfying $g_1(0, 0, 0, y_2, 0) = 0$, or, equivalently, equilibrium points of (31)–(32) on the fold line; that is, satisfying $x_1 = 0$. The type of folded singularity is determined by the Jacobian of

$$\begin{pmatrix} g_{1,x_1}(0, 0, 0, y_2) & g_{1,y_2}(0, 0, 0, y_2) \\ 2g_2(0, 0, 0, y_2) & 0 \end{pmatrix}. \quad (33)$$

Folded saddle nodes (FSN I and FSN II) can arise in the context of systems with two slow and two fast dimensions in an analogous way as described in section 2.2 for systems with two slow and one fast dimensions, and, in the same manner, correspond to the onset of MMOs.

Remark 1. We want to emphasize here the link between the background presented in section 2.1, which deals with 3d system and the computations done above, for the 4D system. Since we have assumed in (23) that $\lambda(y_2) < 0$, we are able to write x_2 as a function of $O(x_1^2)$ by using the implicit function theorem in (27)–(28). By this way, we can obtain a desingularized system in the 4D case that is analogue to the one obtained in the 3D case. This operation can be made in the same way in a system with N fast variables if we assume sufficiently hypothesis on eigenvalues of the Jacobian matrix of the fast subsystem on the fold line.

3. Coupled oscillator system

3.1. Introduction of the system

We consider a system of coupled oscillators in the following form:

$$\varepsilon \dot{x}_1 = -y_1 + F_1(x_1) + \nu H_1^f(x, y) \tag{34}$$

$$\varepsilon \dot{x}_2 = -y_2 + F_2(x_2) + \nu H_2^f(x, y) \tag{35}$$

$$\dot{y}_1 = G_1(x_1, y_1, c_1) + \nu H_1^s(x, y) \tag{36}$$

$$\dot{y}_2 = G_2(x_2, y_2, c_2) + \nu H_2^s(x, y) \quad x = (x_1, x_2) \in \mathbb{R}^2, \quad y = (y_1, y_2) \in \mathbb{R}^2. \tag{37}$$

The parameters ε and ν are the singular parameter and the coupling parameter, respectively, and are considered to be small. The parameters c_1, c_2 control the state of the uncoupled oscillators (moves the nullclines). The coupling functions $H^f : \mathbb{R}^4 \rightarrow \mathbb{R}^2, H^s : \mathbb{R}^4 \rightarrow \mathbb{R}^2$ and $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ are defined by $H^f = (H_1^f, H_2^f)$, by $H^s = (H_1^s, H_2^s)$ and $F = (F_1, F_2)$, respectively. We assume that $y_j = F_j(x)$ are S shaped curves. Written on the fast time scale (34)–(37) has the form:

$$x'_1 = -y_1 + F_1(x_1) + \nu H_1^f(x, y) \tag{38}$$

$$x'_2 = -y_2 + F_2(x_2) + \nu H_2^f(x, y) \tag{39}$$

$$y'_1 = \varepsilon G_1(x_1, y_1, c_1) + \varepsilon \nu H_1^s(x, y) \tag{40}$$

$$y'_2 = \varepsilon G_2(x_2, y_2, c_2) + \varepsilon \nu H_2^s(x, y). \tag{41}$$

We now find the conditions for the existence of a simple fold curve, as in section 2.3. Let Φ be defined as in section 2.3. First note that the critical manifold of (34)–(37), is defined by

$$-y_j + F_j(x) + \nu H_j^f(x, y) = 0, \quad j = 1, 2. \tag{42}$$

The linearization of the rhs of (34)–(35) is given by

$$\begin{pmatrix} F'_1(x_1) + \nu H_{1,x_1}^f & \nu H_{1,x_2}^f \\ \nu H_{2,x_1}^f & F'_2(x_2) + \nu H_{2,x_2}^f \end{pmatrix}, \tag{43}$$

where we assume that (x, y) is on the critical manifold; that is, satisfies (42). The determinant of the matrix in (43) is as follows:

$$F'_1(x_1)F'_2(x_2) + \nu(F'_1(x_1)H_{2,x_2}^f + F'_2(x_2)H_{1,x_1}^f) + \nu^2 \det H_x^f. \tag{44}$$

Proposition 2. Let (x_0, y_0) satisfy $\Phi(x_0, y_0) = 0$ for $\nu = 0$. We assume that

$$F'_1(x_{0,1}) = 0 \quad \text{and} \quad F'_2(x_{0,2}) < 0 \quad \text{as already specified in (23),}$$

and that,

$$F''_1(x_{0,1}) \neq 0.$$

Then, there exists a neighbourhood $U = V \times W$ of $(x_0, y_0) \in \mathbb{R}^4$, and a unique function $\psi : V \subset \mathbb{R} \rightarrow \mathbb{R}^3$, such that:

$$\{(x, y) \in U; \Phi = 0\} = \{(x, y_1) \in \mathbb{R}^3; (x, y_1) = \psi(y_2), y_2 \in V\},$$

with $\psi(y_{2,0}) = (x_{1,0}, x_{2,0}, y_{1,0})$. In addition, the parametrization of the fold curve depends smoothly on the parameters v .

Proof. We can apply proposition 1 but we give here a direct proof by application of the implicit function theorem. Let $M = \Phi_{(x_1, x_2, y_1)}|_{x=0, y=0, v=0}$; then, $\det M = F''(x_{10})(F'(x_{20}))^2 \neq 0$. Note that, because the fast system does not depend on parameters c_1, c_2 , neither does the fold curve.

3.2. Folded singularities and their nature

Let $x_1^* = \psi_1(y_2, v), x_2^* = \psi_2(y_2, v)$ and $y_1^* = \psi_3(y_2, v)$. We transform the fold curve to the line $(0, 0, 0, y_2)$, using the following change of variables: $\tilde{x}_1 = x_1 - x_1^*, \tilde{x}_2 = x_2 - x_2^*, \tilde{y}_1 = y_1 - y_1^*$. We omit the tilde to simplify the notation. System (34)–(37) becomes

$$\varepsilon \dot{x}_1 = -y_1 + f_1(x_1, y_2) + v h_1^f(x, y) + O(\varepsilon) \tag{45}$$

$$\varepsilon \dot{x}_2 = f_2(x_2, y_2) + v h_2^f(x, y) + O(\varepsilon) \tag{46}$$

$$\begin{aligned} \dot{y}_1 &= G_1(x_1 + x_1^*, y_1 + y_1^*, c_1) + v H_1^s(x + x^*, y_1 + y_1^*, y_2) \\ &\quad - \frac{dy_1^*}{dy_2}(G_2(x_2 + x_2^*, y_2, c_2) + v H_2^s(x + x^*, y_1 + y_1^*, y_2)) \end{aligned} \tag{47}$$

$$\dot{y}_2 = G_2(x_2 + x_2^*, y_2, c_2) + v H_2^s(x + x^*, y_1 + y_1^*, y_2), \tag{48}$$

where,

$$f_1(x_1, y_2) = F_1'(x_1^*)x_1 + F_1''(x_1^*)\frac{1}{2}x_1^2 + O(x_1^3),$$

$$f_2(x_2, y_2) = F_2'(x_2^*)x_2 + F_2''(x_2^*)\frac{1}{2}x_2^2 + O(x_2^3),$$

and,

$$h_1^f(x, y) = H_{1,x_1}^f(x^*, y_1^*, y_2)x_1 + H_{1,x_2}^f(x^*, y_1^*, y_2)x_2 + H_{1,y_1}^f(x^*, y_1^*, y_2)y_1 + O(\|(x, y_1)\|^2)$$

$$h_2^f(x, y) = H_{2,x_1}^f(x^*, y_1^*, y_2)x_1 + H_{2,x_2}^f(x^*, y_1^*, y_2)x_2 + H_{2,y_1}^f(x^*, y_1^*, y_2)y_1 + O(\|(x, y_1)\|^2).$$

Our goal is to arrive at the desingularized system (31)–(32). The first step is to diagonalize the linear part of the fast subsystem (45)–(46). More precisely, we change coordinates so that the rhs of (45)–(48) is transformed to the form (26). Let

$$\begin{pmatrix} 1 \\ v(y_2) \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} w(y_2) \\ 1 \end{pmatrix} \tag{49}$$

be the eigenvectors of the Jacobian (43) at the points $(0, 0, 0, y_2)$ on the fold line, corresponding to the eigenvalues 0 and $\lambda(y_2) < 0$, respectively. The diagonalizing transformation has the form

$$\tilde{x} = P(y)^{-1}x,$$

where the columns of the matrix P are the eigenvectors (49). In the new variables system (45)–(48) becomes (tilde is omitted)

$$\varepsilon \dot{x}_1 = \frac{1}{1 - vw} \left(-y_1 + \frac{K}{2} x_1^2 + v(O(y_1)) + O(x_1 x_2, x_2^2) + O(\varepsilon) \right) \tag{50}$$

$$\varepsilon \dot{x}_2 = \lambda(y_2)x_2 + v(O(y_1)) + O(\|x\|^2) + O(\varepsilon) \tag{51}$$

$$\begin{aligned} \dot{y}_1 &= G_1(x_1 + x_2 w(y_2) + x_1^*, y_1 + y_1^*, c_1) + vH_1^s(P(y_2)x + x^*, y_1 + y_1^*, y_2) \\ &\quad - \frac{dy_1^*}{dy_2} \left(G_2(x_2 + x_1 v(y_2) + x_2^*, y_2, c_2) + vH_2^s(P(y_2)x + x^*, y_1 + y_1^*, y_2) \right) \end{aligned} \tag{52}$$

$$\dot{y}_2 = G_2(x_2 + x_1 v(y_2) + x_2^*, y_2, c_2) + vH_2^s(P(y_2)x + x^*, y_1 + y_1^*, y_2), \tag{53}$$

where,

$$K = F_1''(x_1^*) + O(v). \tag{54}$$

WLOG, in the remaining of this section, we will assume that $F_1''(x_1^*) > 0$. Note that the fast subsystem (50)–(51) is now as specified in (26). By applying the procedure described in section 2.3 we obtain the reduced system

$$(Kx_1 + O(x_1^2))\dot{x}_1 = g_1(x_1, y_2, c_1, c_2), \tag{55}$$

$$\dot{y}_2 = g_2(x_1, y_2, c_2), \tag{56}$$

where,

$$\begin{aligned} g_1(x_1, y_2, c_1, c_2) &= G_1(x_1^* + x_1 + w(y_2)O(x_1^2), y_1^* + O(x_1^2), c_1) \\ &\quad + vH_1^s(x_1^* + x_1 + w(y_2)O(x_1^2), x_2^* + v(y_2)x_1 + O(x_1^2), y_1^* + O(x_1^2), y_2) \\ &\quad - \frac{dy_1^*}{dy_2} \left(G_2(x_2^* + v(y_2)x_1 + O(x_1^2)) + vH_2^s(x_1^* + x_1 + w(y_2)O(x_1^2), \right. \\ &\quad \left. x_2^* + v(y_2)x_1 + O(x_1^2), y_1^* + O(x_1^2), y_2) \right) \end{aligned} \tag{57}$$

$$\begin{aligned} g_2(x_1, y_2, c_2) &= G_2(x_2^* + v(y_2)x_1 + O(x_1^2)) + vH_2^s(x_1^* + x_1 + w(y_2)O(x_1^2), \\ &\quad x_2^* + v(y_2)x_1 + O(x_1^2), y_1^* + O(x_1^2), y_2), \end{aligned} \tag{58}$$

and the desingularized system

$$\dot{x}_1 = g_1(x_1, y_2, c_1, c_2), \tag{59}$$

$$\dot{y}_2 = g_2(x_1, y_2, c_2)(2Kx_1 + O(x_1^2)). \tag{60}$$

3.3. Folded singularities of type FSN II and the existence of MMOs

As discussed in section 2.2, a well known mechanism of transition to MMOs is FSN II, see [9, 16]. This is a codimension one transition, corresponding to the passage of the system from a parameter region with an excitable equilibrium to a parameter region of MMO dynamics. This can be described as follows in the context of system (55)–(56): as the regular parameter is varied a stable equilibrium of (55)–(56) approaches the fold, and, for the critical value of the regular parameter, satisfies the conditions:

$$g_1(0, y_2, c_1, c_2) = 0 \tag{61}$$

$$g_2(0, y_2, c_2) = 0. \tag{62}$$

On the other side of criticality there are MMOs as well as an equilibrium, which are now unstable.

In the following, we assume that the parameter c_2 is fixed. Note that for $v = 0$, y_1^* does not depend on y_2 and thus $dy_1^*/dy_2 = O(v)$. We assume that, for $v = 0$, the equation




Figure 1. The $\nu = 0$ limit configurations. (a) The uncoupled system (38)–(40) admits a unique repulsive stationary point near the fold. (b) The uncoupled system (39)–(41) admits a unique attractive stationary point on the stable manifold.

$G_2(x_2^*, y_2, c_2) = 0$ has a unique solution $x_2^* = \psi_2(y_2, c_2)$, which is a stable equilibrium of the uncoupled (x_2, y_2) subsystem, and that $G_{2,y_2} \neq 0$ at this point. By the implicit function theorem, this gives a unique solution y_2 for the equation (62), for ν small enough. Recall that, by hypothesis, in the case $\nu = 0$ we assume that, each uncoupled subsystem (38)–(40) and (39)–(41) admits a unique stationary point, one attractive on the stable manifold and one repulsive near the fold. This corresponds to the nullcline configuration shown in figure 1. Let us denote this particular value \bar{y}_2 . It follows that this value \bar{y}_2 determines particular values of $(x_1^*, y_1^*, x_2^*) = (\bar{x}_1^*, \bar{y}_1^*, \bar{x}_2^*)$. We assume that solving equation (61) with these values gives a unique value of $c_1 = \bar{c}_1$.

Recall from the discussion in section 2.2 that a non-degenerate FSN II singularity corresponds to a transcritical bifurcation for system (59)–(60).

The following proposition establishes the existence of a transcritical bifurcation for (59)–(60) and the existence of FSN II.

Proposition 3. *Let us assume that, for system (59)–(60):*

$$\frac{dg_1}{dy_2} \neq 0 \quad \text{and} \quad \frac{dG_1}{dx_1} \frac{dG_2}{dy_2} \neq 0. \tag{63}$$

Then, for c_1 in a neighbourhood of $c_1 = \bar{c}_1$ and ν small enough, there exists two stationary points for system (59)–(60): $(x_{1,e}, y_{2,e})$ and $(0, y_{2,\text{fold}})$. Furthermore, if we assume that

$$\frac{dg_1}{dy_2} \frac{dg_2(0, y_{2,\text{fold}}(c_1))}{dc_1} < 0, \quad G_{1,x_1} > 0 \quad \text{and} \quad G_{2,y_2} \frac{dx_{1,e}}{dc_1} < 0, \tag{64}$$

then, for ν sufficiently small, there is a transcritical bifurcation. As the parameter c_1 increases from left to right, the folded stationary point passes from a folded saddle to a repulsive folded node, whereas the ordinary stationary point passes from a repulsive node to a saddle. It follows that $(0, \bar{y}_2)$, is an FSN II point.

Proof. The folded stationary point, $(0, y_{2,\text{fold}})$ is obtained by solving the equation $g_1(0, y_2, c_1, c_2) = 0$, i.e., $g_1 = 0, x_1 = 0$. For this, we apply the implicit function theorem to $g_1(0, y_2, c_1, c_2)$ as a function of y_2 and c_1 . For $y_2 = \bar{y}_2$ and $c_1 = \bar{c}_1$, we have $g_1 = 0$. From hypothesis (63), it follows that $\frac{dg_1}{dy_2}(0, \bar{y}_2, \bar{c}_1, c_2) \neq 0$, then the implicit function theorem gives the existence of a stationary point $(x_1 = 0, y_2 = y_{2,\text{fold}})$ for c_1 in

a neighbourhood of \bar{c}_1 . The second stationary point, $(x_{1,e}, y_{2,e})$, is obtained by solving the equation $g_1(x_1, y_2, c_1) = 0, g_2(x_1, y_2) = 0$. Note that this second stationary point corresponds to a true stationary point of system (34)–(37). For $\nu = 0$, we know that these equations admit a unique solution. Using hypothesis (64), we can apply the implicit function theorem in a neighbourhood of $\nu = 0$. This gives the existence of a unique stationary point for ν sufficiently small. For the first stationary point, the folded one $(0, y_{2,\text{fold}})$ the Jacobian is given by

$$J_1 = \begin{pmatrix} G_{1,x_1} + O(\nu) & \frac{dg_1}{dy_2} \\ 2Kg_2 & 0 \end{pmatrix}. \tag{65}$$

This gives:

$$\det J_1 = - \left(\frac{dg_1}{dy_2} \right) 2Kg_2.$$

For the second stationary point, the ordinary one, the Jacobian is given by:

$$J_2 = \begin{pmatrix} G_{1,x_1} + O(\nu) + O(x_1) & \frac{dg_1}{dy_2} \\ KG_{2,x_1}x_1 + O(x_1^2) & KG_{2,y_2}x_1 + O(x_1^2) \end{pmatrix}. \tag{66}$$

This gives:

$$\det J_2 = KG_{1,x_1}G_{2,y_2}x_1 + O(x_1\nu).$$

From hypothesis (64), it follows that, as c_1 crosses the value \bar{c}_1 , $\det J_1$ crosses the value 0 from negative to positive whereas $\det J_2$ crosses the value 0 from positive to negative. Subsequently the stationary point $(0, y_{2,\text{fold}})$ bifurcates from a saddle to a repulsive node whereas the stationary point $(x_{1,e}, y_{2,e})$ bifurcates from a repulsive node to a saddle. Then, it follows that the system (59)–(60) admits a transcritical bifurcation at point $(0, \bar{y}_2)$ when c_1 crosses the value \bar{c}_1 , and $(0, \bar{y}_2)$ is a FSN II.

4. Example—coupled FitzHugh–Nagumo systems

4.1. Simple fold line

We consider the following system

$$\begin{cases} \epsilon \dot{x}_1 = F(x_1) - y_1 + \alpha_1(x_2 - x_1) \\ \epsilon \dot{x}_2 = F(x_2) - y_2 + \alpha_2(x_1 - x_2) \\ \dot{y}_1 = x_1 - c_1 + \beta_1(y_2 - y_1) \\ \dot{y}_2 = x_2 - c_2 + \beta_2(y_1 - y_2) \end{cases} \tag{67}$$

with

$$F(z) = -z^3 + 3z.$$

We define

$$l(x_1, y_1, x_2, y_2) = \begin{cases} F(x_1) - y_1 + \alpha_1(x_2 - x_1) \\ F(x_2) - y_2 + \alpha_2(x_1 - x_2). \end{cases} \tag{68}$$

Points on the fold line are defined by the condition

$$\begin{cases} \det(Dl(x, y)) = 0 \\ l_1(x, y) = 0 \\ l_2(x, y) = 0 \end{cases} \tag{69}$$

which, for system (67), is given by

$$\begin{cases} (F'(x_1) - \alpha_1)(F'(x_2) - \alpha_2) - \alpha_1\alpha_2 = 0 \\ y_1 = F(x_1) + \alpha_1(x_2 - x_1) \\ y_2 = F(x_2) + \alpha_2(x_1 - x_2). \end{cases}$$

Thanks to proposition 2, the existence of a *simple* fold line follows from $F'(x_1) = 0$, $F'(x_2) \neq 0$ and $F''(x_1) \neq 0$ (the alternative choice would be $F'(x_2) = 0$, $F'(x_1) \neq 0$ and $F''(x_2) \neq 0$). We obtain a solution of system (4.1) with (x_1, x_2, y_1) given as function of y_2 . We denote this solution by $(x_1^*, x_2^*, y_1^*) = (\varphi_1(y_2), \varphi_2(y_2), \varphi_3(y_2))$. Note that we have obtained a curve in \mathbb{R}^4 parametrized by y_2 , see section 2 for a general statement.

4.2. Folded singularities and their nature

System (45)–(48) in context of (67) becomes

$$\begin{cases} \epsilon \dot{x}_1 = -y_1 + f_1(x_1, y_2) + \alpha_1(x_2 - x_1) + O(\epsilon) \\ \epsilon \dot{x}_2 = f_2(x_2, y_2) + \alpha_2(x_1 - x_2) + O(\epsilon) \\ \dot{y}_1 = x_1^* + x_1 - c_1 + \beta_1(y_2 - (y_1^* + y_1)) \\ \quad - \frac{dy_1^*}{dy_2}(x_2^* + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2)) \\ \dot{y}_2 = x_2^* + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2) \end{cases} \tag{70}$$

where

$$f_1(x_1) = F'(x_1^*)x_1 + F''(x_1^*)\frac{x_1^2}{2} - x_1^3$$

and

$$f_2(x_2) = F'(x_2^*)x_2 + F''(x_2^*)\frac{x_2^2}{2} - x_2^3.$$

As in section 3.2, we diagonalize the linear part of the fast system. The eigenvalues of $Df(x_1^*, x_2^*)$ are:

$$0 \quad \text{and} \quad \lambda(y_2) = F'(x_1^*) + F'(x_2^*) - \alpha_1 - \alpha_2.$$

As explained in section 3.3, we focus on the dynamics described by configuration in figure 1, thus $F'(x_2^*) < 0$. The associated eigenvectors are, respectively,

$$P_1 = \begin{pmatrix} 1 \\ \frac{\alpha_2}{\alpha_2 - F'(x_2^*)} \end{pmatrix} \quad \text{and} \quad P_2 = \begin{pmatrix} \frac{\alpha_1}{F'(x_1^*) - \alpha_2} \\ 1 \end{pmatrix}.$$

We now diagonalize the fast subsystem using the transformation

$$X = P\tilde{X} \quad \text{with} \quad X = \begin{pmatrix} x_1 \\ x_2 \end{pmatrix}$$

where P is the matrix whose columns are the eigenvectors. We introduce the following notations:

$$w = \frac{\alpha_1}{F'(x_2^*) - \alpha_2}, \quad v = \frac{\alpha_2}{\alpha_2 - F'(x_2^*)} \quad d = \frac{1}{1 - vw}.$$

After some transformations , system (70) becomes:

$$\begin{cases} \epsilon \dot{x}_1 = d((F''(x_1^*) \frac{(x_1 + wx_2)^2}{2} - (x_1 + wx_2)^3 - y_1) \\ \quad - w(F''(x_2^*) \frac{(vx_1 + x_2)^2}{2} - (vx_1 + x_2)^3)) + O(\epsilon) \\ \epsilon \dot{x}_2 = \lambda(y_2)x_2 + d(-v(F''(x_1^*) \frac{(x_1 + wx_2)^2}{2} - (x_1 + wx_2)^3 - y_1) \\ \quad + F''(x_2^*) \frac{(vx_1 + x_2)^2}{2} - (vx_1 + x_2)^3) + O(\epsilon) \\ \dot{y}_1 = x_1^* + x_1 + wx_2 - c_1 + \beta_1(y_2 - (y_1^* + y_1)) \\ \quad - \frac{dy_1^*}{dy_2}(x_2^* + vx_1 + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2)) \\ \dot{y}_2 = x_2^* + vx_1 + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2). \end{cases}$$

Hence, the reduced equation is given by

$$0 = d((F''(x_1^*) \frac{(x_1 + wx_2)^2}{2} - (x_1 + wx_2)^3 - y_1) - w(F''(x_2^*) \frac{(vx_1 + x_2)^2}{2} - (vx_1 + x_2)^3)) \tag{71}$$

$$0 = \lambda(y_2)x_2 + d(-v(F''(x_1^*) \frac{(x_1 + wx_2)^2}{2} - (x_1 + wx_2)^3 - y_1) + F''(x_2^*) \frac{(vx_1 + x_2)^2}{2} - (vx_1 + x_2)^3) \tag{72}$$

$$\dot{y}_1 = x_1^* + x_1 + wx_2 - c_1 + \beta_1(y_2 - (y_1^* + y_1)) - \frac{dy_1^*}{dy_2}(x_2^* + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2)) \tag{73}$$

$$\dot{y}_2 = x_2^* + vx_1 + x_2 - c_2 + \beta_2(y_1^* + y_1 - y_2). \tag{74}$$

Now, using equation (72) and the implicit function theorem in (73), as $\lambda(y_2) < 0$, one can obtain x_2 as a function of x_1, y_2 wich leads to,

$$x_2 = O(x_1^2).$$

Now, (72) can be rewritten in the form

$$y_1 = \frac{K}{2}x_1^2 + O(x_1^3) \tag{75}$$

with

$$\begin{aligned} K &= F''(x_1^*) - wv^2 F''(x_2^*) \\ &= F''(x_1^*) + O(v^3), \end{aligned}$$

where $0 \leq v \leq \max(\alpha_1, \beta_1, \alpha_2, \beta_2)$. Subsequently, we derivate (75), and plug into (73) and (74). We obtain,

$$\begin{aligned} (Kx_1 + O(x_1^2))\dot{x}_1 &= x_1^* + x_1 - c_1 + \beta_1(y_2 - y_1^*) \\ &\quad - \frac{dy_1^*}{dy_2}(x_2^* + bx_1 - c_2 + \beta_2(y_1^* - y_2)) + O(x_1^2) \\ \dot{y}_2 &= x_2^* + bx_1 - c_2 + \beta_2(y_1^* - y_2) + O(x_1^2). \end{aligned}$$

This gives the following desingularized system:

$$\dot{x}_1 = x_1^* + x_1 - c_1 + \beta_1(y_2 - y_1^*) - \frac{dy_1^*}{dy_2}(x_2^* + vx_1 - c_2 + \beta_2(y_1^* - y_2)) + O(x_1^2) \tag{76}$$

$$\dot{y}_2 = (x_2^* + vx_1 - c_2 + \beta_2(y_1^* - y_2) + O(x_1^2))(Kx_1 + O(x_1^2)). \tag{77}$$

$$\tag{78}$$

Hence, using notations of section 3.3, we have:

$$g_1(x_1, y_2) = x_1^* + x_1 - c_1 + \beta_1(y_2 - y_1^*) - \frac{\partial y_1^*}{\partial y_2}(x_2^* + vx_1 - c_2 + \beta_2(y_1^* - y_2)) + O(x_1^2)$$

and

$$g_2(x_1, y_2) = x_2^* + vx_1 - c_2 + \beta_2(y_1^* - y_2) + O(x_1^2).$$

Proposition 4. *Let us assume $c_2 < -1$ and $\beta_1 - \frac{\alpha_1}{9(1-c_2^2)^2} > 0$. For $\alpha_1, \alpha_2, \beta_1, \beta_2$ small enough, let \bar{y}_2 be the solution of $g_2(0, y_2) = 0$ and define $\bar{c}_1 = x_1^* + \beta_1(\bar{y}_2 - y_1^*)$. Then, for $\alpha_1, \alpha_2, \beta_1, \beta_2$ in a small neighbourhood of zero, the system (76)–(77), admits a transcritical bifurcation at point $x_1 = 0, y_2 = \bar{y}_2, c_1 = \bar{c}_1$. As the parameter c_1 crosses the value \bar{c}_1 from left to right, the folded stationary point passes from a folded saddle to a repulsive folded node, whereas the ordinary stationary point passes from a repulsive node to a saddle.*

Proof. This proposition is a specific case of proposition (3). We will verify that:

$$\frac{dg_1}{dy_2}(0, \bar{y}_2, \bar{c}_1) > 0, \tag{79}$$

$$\frac{dG_1}{dx_1} > 0, \tag{80}$$

$$\frac{dG_2}{dy_2} < 0, \tag{81}$$

$$\frac{dx_{1,e}}{dc_1} > 0, \tag{82}$$

$$\frac{dg_2(0, \bar{y}_{2, \text{fold}}(c_1))}{dc_1} < 0. \tag{83}$$

We start with hypothesis (79). After some computation, we find that this hypothesis reads as,

$$\beta_1 - \frac{\alpha_1}{9(1 - \bar{x}_2^{*2})^2} + O(v^2) > 0.$$

This holds since $\bar{x}_2^* = c_2 + O(v)$. Hypothesis (80), is verified since $\frac{\partial G_1}{\partial x_1} = 1$.

Now, we deal with (81). Here, it reads:

$$\frac{dx_2^*}{dy_2} < 0. \tag{84}$$

This holds since $\frac{dx_2^*}{dy_2} = \frac{1}{F'(x_2)} + O(v) < 0$ for v sufficiently small. Hypothesis (82) is verified since we have $\frac{dx_{1,e}}{dc_1} = 1 + O(v) > 0$.

Finally, we come to hypothesis (83). From (79), we know that $\frac{dg_1}{dy_2}(0, y_2) > 0$. It follows that $\frac{y_{2, \text{fold}}}{dc_1} > 0$. Furthermore, we have that $\frac{dg_2}{dy_2}(0, y_2) < 0$. We conclude that $\frac{dg_2(0, \bar{y}_{2, \text{fold}}(c_1))}{dc_1} < 0$.




Figure 2. This figure corresponds to the simulation of system (67) for a value of parameter $c_1 = -0.953$. In (a) we show the time evolution of x_1 in the interval $[0, 500]$. In (b) we show the behaviour of the system in the (x_1, y_1) phase-plane. For this value of the parameter c_1 the system evolves asymptotically to this stationary state.

4.3. Numerical simulations

We have performed numerical integration of system (67) on time interval $[0, 500]$, using a fourth-order Runge–Kutta method with time step of 0.0001. The parameter values are the following: $\varepsilon = 0.01, \alpha_1 = \beta_1 = \alpha_2 = \beta_2 = 0.05, c_2 = -1.5$. As explained in proposition 4, we obtain the critical value of parameter \bar{c}_1 by solving the following equations:

$$g_2(0, y_2) = 0$$

$$c_1 = x_1^* + \beta_1(y_2 - y_1^*).$$

In our case, we find:

$$\bar{c}_1 \simeq -0.95266.$$

Then, we vary the parameter c_1 in a small neighbourhood around the value of \bar{c}_1 . If $c_1 < \bar{c}_1$, then the system goes to a stationary state. As c_1 crosses the value \bar{c}_1 , we can observe the appearance of MMOs. Below, we illustrate the simulation of system (67) for two values of c_1 : for $c_1 = -0.953$, which corresponds to a case where the system goes to the stationary state, and for $c_1 = -0.952$, which corresponds to a case where we observe MMOs. Furthermore, we can approximate, for the folded stationary point, the eigenvalues of the Jacobian of (76)–(77). For $c_1 = -0.953$, we find:

$$\lambda_1 \simeq 0.998, \quad \lambda_2 \simeq -0.02.$$

And for $c_1 = -0.952$, we find:

$$\lambda_1 \simeq 1.004, \quad \lambda_2 \simeq 0.004.$$

Following [3, 16], this gives an approximate theoretical number of small oscillations:

$$s = \left\lceil \frac{1}{2} + \frac{\lambda_1}{\lambda_2} \right\rceil \simeq 135.$$

In figure 2, we have illustrated the results of the simulation, for the parameter value $c_1 = -0.953$. It shows that in this case the system evolves asymptotically towards the stationary




Figure 3. This figure corresponds to the simulation of system (67) for a value of parameter $c_1 = -0.952$. In (a) we show the time evolution of x_1 , for t in the interval $[400,450]$. In (b) we show the behaviour of the system in the (x_1, y_1) phase-plane. For this value of the parameter c_1 , the MMOs appear. But the amplitude of the small oscillations is very small compared to that of the large oscillations, and a zoom in is necessary to distinguish them.




Figure 4. This figure corresponds to the simulation of system (67) for a value of parameter $c_1 = -0.952$. It is the same simulation as the one represented in figure 3, but with a zoom on the zone of small oscillations. In (a) we show the time evolution of x_1 , for t in the interval $[400,450]$ and x_1 in the interval $[-1.1, -0.9]$. In (b) we show the behaviour of the system in the (x_1, y_1) phase-plane, for x_1 in the interval $[-1.1, -0.9]$ and y_1 in the interval $[-2.05, -2]$. We can see, the trajectory on the attractive manifold, then the small oscillations, resulting from the intersection of attractive and repulsive manifolds, and finally, the trajectory along the repulsive manifold before the evolution towards the fast direction.

state. The other figures illustrate the simulation results for $c_1 = -0.952$. In figure 3(a), we show the time evolution of the fast variable x_1 , whereas in figure 3(b) we show the behaviour in the (x_1, y_1) phase-plane. Here, the MMOs appear but the small oscillations are hardly distinguishable because of their tiny amplitude in comparison with the relaxation oscillations. So, in figures 4 and 5 we show a zoom of the previous illustration. In figure 4(a), we show the time evolution of the fast variable x_1 , whereas in figure 4(b), we show the behaviour in the




Figure 5. This figure corresponds to the simulation of system (67) for a value of parameter $c_1 = -0.952$. It is the same simulation as the one represented in figure 3, but with a zoom on the area of small oscillations. We show the behaviour of the system in the (x_1, y_2, y_1) phase-plane, for x_1 in the interval $[-1.1, -0.9]$ and y_1 in the interval $[-2.026, -2.016]$. Here, we can easily distinguish the behaviour described below. The small oscillations occur when the trajectories originating in the attracting manifold in one side of the strong canard are trapped by the repelling one and have to return to the attracting manifold. On the other side of the folded node, the trajectories follow the fast direction, see [3].




Figure 6. This figure shows the simulation of system (67) for parameter $c_1 = -0.952$. We can see the critical manifold defined by equations: $(x_1, y_2 = F(x_2) + \alpha_2(x_1 - x_2), y_1 = F(x_1) + \alpha_1(x_2 - x_1))$ and the behaviour of the system in the (x_1, y_2, y_1) phase-space (green curve). It shows the trajectory along the attractive critical manifold, then the evolution near the fold curve, the evolution along the fast direction and the global return mechanism.

(x_1, y_1) phase-plane. In panel (a) we can easily distinguish the small oscillations. In panel (b) we can see the trajectory on the attractive manifold, then the small oscillations, resulting from the intersection of attractive and repulsive manifolds, and finally, the trajectory along the repulsive manifold before the evolution toward the fast direction. Figure 5 is interesting as it clearly illustrates, in the (x_1, y_2, y_1) phase plane, the MMOs in the case of folded node singularity. The trajectories of system close to the singular funnel enter a region near the fold where they rotate around the weak canard: the trajectories originating in the attracting manifold in one side of the strong canard are trapped by the repelling one and have to return




Figure 7. This figure shows the simulation of system (67) for parameter $c_1 = -0.952$. We can see the critical manifold defined by equations: $(x_1, y_2 = F(x_2) + \alpha_2(x_1 - x_2), y_1 = F(x_1) + \alpha_1(x_2 - x_1))$ and the behaviour of the system in the (x_1, y_2, y_1) phase-space (green curve). It shows the trajectory along the attractive critical manifold, then the evolution near the fold curve and finally the evolution along the fast direction.




Figure 8. This figure shows the simulation of system (67) for parameter $c_1 = -0.952$. We can see the critical manifold defined by equations: $(x_1, y_2 = F(x_2) + \alpha_2(x_1 - x_2), y_1 = F(x_1) + \alpha_1(x_2 - x_1))$ and the behaviour of the system in the (x_1, y_2, y_1) phase-space (green curve). It shows the small oscillations occurring when the trajectory oscillates around the weak canard. The amplitude of oscillations when entering in the funnel region start to decrease before increasing until they cross the folded node and follow the fast direction.

to the attracting manifold, see [3] for further detail. Finally, figures 6, 7 and 8, also show the MMOs in the space (x_1, y_2, y_1) . In figure 6, we show the critical manifold defined by $(x_1, y_2 = F(x_2) + \alpha_2(x_1 - x_2), y_1 = F(x_1) + \alpha_1(x_2 - x_1))$, and the trajectory of the system in the (x_1, y_2, y_1) phase-space. We can see the small oscillations near the fold, the trajectory along the fast direction and the global return mechanism. The figure 7 is the same as the figure 6, but with a zoom on area of the fold. The same is done, in figure 8 but with a greater zoom in the area of small oscillations.

5. An alternative approach to the analysis of coupled oscillator systems

In this section we outline an alternative approach that was pointed out to us by an anonymous referee. The modification consists of defining the reduced flow of (67) by means of the fast variables x_1 and x_2 . Differentiating the equation $l(x_1, x_2, y_1, y_2) = 0$ (see (68) for the definition) we obtain

$$\begin{pmatrix} \dot{y}_1 \\ \dot{y}_2 \end{pmatrix} = \begin{pmatrix} F'(x_1) - \alpha_1 & \alpha_1 \\ \alpha_2 & F'(x_2) - \alpha_2 \end{pmatrix} \begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \end{pmatrix}. \tag{85}$$

Hence, the reduced equation can be written in the form:

$$\begin{aligned} & \begin{pmatrix} F'(x_1) - \alpha_1 & \alpha_1 \\ \alpha_2 & F'(x_2) - \alpha_2 \end{pmatrix} \begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \end{pmatrix} \\ &= \begin{pmatrix} x_1 - c_1 + \beta_1(F(x_2) - F(x_1) - (\alpha_1 + \alpha_2)(x_2 - x_1)) \\ x_2 - c_2 + \beta_2(F(x_2) - F(x_1) - (\alpha_1 + \alpha_2)(x_2 - x_1)) \end{pmatrix}. \end{aligned} \tag{86}$$

We multiply both sides of (86) by the cofactor matrix

$$\begin{pmatrix} F'(x_2) - \alpha_2 & -\alpha_1 \\ -\alpha_2 & F'(x_1) - \alpha_1 \end{pmatrix}, \tag{87}$$

which yields

$$\begin{aligned} & ((F'(x_1) - \alpha_1)(F'(x_2) - \alpha_2) - \alpha_1\alpha_2) \begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \end{pmatrix} \\ &= \begin{pmatrix} F'(x_2) - \alpha_2 & -\alpha_1 \\ -\alpha_2 & F'(x_1) - \alpha_1 \end{pmatrix} \begin{pmatrix} x_1 - c_1 + \beta_1(F(x_2) - F(x_1) - (\alpha_1 + \alpha_2)(x_2 - x_1)) \\ x_2 - c_2 + \beta_2(F(x_2) - F(x_1) - (\alpha_1 + \alpha_2)(x_2 - x_1)) \end{pmatrix}. \end{aligned} \tag{88}$$

Now we can desingularize by cancelling the factor $(F'(x_1) - \alpha_1)(F'(x_2) - \alpha_2) - \alpha_1\alpha_2$, which results in the following desingularized system

$$\begin{aligned} & \begin{pmatrix} \dot{x}_1 \\ \dot{x}_2 \end{pmatrix} = \\ & \begin{pmatrix} F'(x_2) - \alpha_2 & -\alpha_1 \\ -\alpha_2 & F'(x_1) - \alpha_1 \end{pmatrix} \begin{pmatrix} x_1 - c_1 + \beta_1(F(x_2) - F(x_1) - (\alpha_1 + \alpha_2)(x_2 - x_1)) \\ x_2 - c_2 + \beta_2(F(x_2) - F(x_1) - (\alpha_1 + \alpha_2)(x_2 - x_1)) \end{pmatrix}. \end{aligned} \tag{89}$$

This is equivalent to the desingularization that takes (7)–(8) to (9)–(10). In the context of (89) folded singularities are equilibria on the fold line defined by $((F'(x_1) - \alpha_1)(F'(x_2) - \alpha_2) - \alpha_1\alpha_2) = 0$. Note that, such equilibria exist robustly since the cofactor matrix is singular along the fold line. To verify that we have a FSN II, we have to find a transcritical bifurcation, with one equilibrium crossing the fold line and the other one staying on it.

This approach seems viable and may have computational advantages over the one we have used. It is suited to the context of coupled oscillators, but relies on the fact that there are at least as many fast variables as there are slow variables.

6. Conclusion

In this paper, we have studied a system of two coupled slow–fast oscillators. We showed that coupling them leads to the occurrence of MMOs. The main observation was that coupling the two systems have rise to an FSN II point, which is known to imply the existence of small amplitude oscillations in the fold region. Our study has been the first attempt to understand canards and MMOs in systems of coupled oscillators. Our future goal is to extend our work to the context of large systems of oscillators. As coupled oscillators systems arise

as discretizations of reaction diffusion equations, we hope to use the insights of this and future work to understand MMOs in the context of reaction diffusion equations. Since numerous models in biology and neuroscience are constructed as networks of oscillators and reaction diffusion equations, this research has to be done with deep interactions with applications.

Acknowledgment

We would like to thank Région Haute Normandie and FEDER (RISC project) for financial support.

References

- [1] Benoît E 1990 Canards et enlacements *Publ. Math. IHES* **72** 63–91
- [2] Benoît E, Callot J-L, Diener F and Diener M 1981 Chasse au canard *Collect. Math.* **31** 37–119
- [3] Brøns M, Krupa M and Wechselberger M 2006 Mixed-mode oscillations due to generalized canard phenomenon *Bifurcation Theory and Spatio-Temporal Pattern Formation* vol 49 (*Fields Institute Communications*) (Providence, RI: American Mathematical Society) pp 39–64
- [4] Brøns M 1988 Bifurcations and instabilities in the Greitzer model for compressor system surge *Math. Eng. Industry* **2** 51–63
- [5] Curtu R and Rubin J 2011 Interaction of canard and singular Hopf mechanisms in a neural model *SIAM J. Appl. Dyn. Syst.* **10** 1443–79
- [6] Desroches M, Guckenheimer J, Krauskopf B, Kuehn C, Osinga H M and Wechselberger M 2012 Mixed-mode oscillations with multiple time scales *SIAM Rev.* **54** 211–88
- [7] Ermentrout B and Wechselberger M 2009 Canards, clusters and synchronization in a weakly coupled interneuron model *SIAM J. Appl. Dyn. Syst.* **8** 253–78
- [8] Guckenheimer J 2008 Singular Hopf bifurcation in systems with two slow variables *SIAM J. Appl. Dyn. Syst.* **7** 1355–77
- [9] Krupa M and Wechselberger M 2010 Local analysis near a folded saddle-node singularity *J. Diff. Eqns* **248** 2841–88
- [10] Milik A, Szmolyan P, Loeffelmann H and Groeller E 1998 Geometry of mixed-mode oscillations in the 3D autocatalator *Int. J. Bifur. Chaos* **8** 505–19
- [11] Rotstein H, Wechselberger M and Kopell N 2008 Canard induced mixed-mode oscillations in a medial entorhinal cortex layer II stellate cell model *SIAM J. Appl. Dyn. Syst.* **7** 1582–611
- [12] Rubin J and Wechselberger M 2007 Giant squid-hidden canard: the 3D geometry of the Hodgkin–Huxley model *Biol. Cybernet.* **97** 5–32
- [13] Szmolyan P and Wechselberger M 2001 Canards in \mathbb{R}^3 *J. Diff. Eqns* **177** 419–53
- [14] Vo T, Bertram R, Tabak J and Wechselberger M 2010 Mixed mode oscillations as a mechanism for pseudo-plateau bursting *J. Comput. Neurosci.* **28** 443–58
- [15] Wechselberger M 2005 Existence and bifurcations of canards in \mathbb{R}^3 in the case of the folded node *SIAM J. Appl. Dyn. Syst.* **4** 101–39
- [16] Wechselberger M 2007 Canards *Scholarpedia* **2** 1356
- [17] Wechselberger M 2011 A propos de canards (apropos canards) *Trans. Am. Math. Soc.* **364** 3289–309